

SiO Observations of the NGC 1333 IRAS 2A Protostellar Jet

F.P. Wilkin (f.wilkin@astrosmo.unam.mx)*
Instituto de Astronomía, UNAM Morelia, México

H.-R. Chen
Astronomy Department, University of California, Berkeley, USA

L.M. Chernin
2694 Barkley Ave, Santa Clara, CA, USA

R.L. Plambeck
Astronomy Department, University of California, Berkeley, USA

Abstract. We present 86.8 GHz SiO J=2-1 observations of the molecular outflow from the protostellar source NGC 1333 IRAS 2A. Silicon monoxide is a sensitive tracer of shocks in molecular gas, and the emission clearly traces terminal bowshocks at the ends of the outflow lobes, as well as emission knots and a source coincident with the protostar. Our three-field mosaic from the BIMA interferometer covers the entire outflow length, including both lobes. The most prominent emission feature, the eastern bowshock, shows SiO at the leading edge, clearly displaced from the maximum seen in CO. This is in contrast to recent observations by Chandler & Richer (2001) of the HH 211 jet, where SiO traced primarily internal working surfaces and was less extended than the CO. Our results are consistent with prompt entrainment at the leading edge of the IRAS 2A bowshocks.

Keywords: Protostars, Molecular Outflows, Shocks

1. Summary

The class 0 source IRAS 2A drives an unusual, highly collimated jet-like outflow first studied in CO (Sandell *et al.* 1994). Such a young, symmetric, and focused outflow provides an excellent opportunity to observe the expected bow shock of a jet interacting with the molecular ambient gas. We used the BIMA interferometer to map this outflow in SiO in order to trace shocks within the outflow. Our initial three-field, C-array ($10'' \times 7''$ beam) mosaic covers the positions of IRAS 2A and the east and west bow shocks. In addition to this outflow of roughly E-W orientation, a larger, less collimated bipolar outflow originates from nearly the same position (Liseau *et al.* 1988), and is possibly associated with the nearby source IRAS 2C (Knee & Sandell 2000; Sandell & Knee 2001). Because the position of IRAS 2C agrees poorly with

* Partially supported by NSF International Researchers Fellows Program



the symmetry axes of the flows seen in CO, the possibility exists that the two perpendicular outflows seen in CO might actually represent a single, much wider, limb-brightened bipolar outflow (see Figure 1). Our recent observations with BIMA in the D-array (13.5'' beam) detect





Figure 1. Mosaic of the IRAS 2A region. Thin contours show 12CO J=2-1 taken from the BIMA observations of Engargiola & Plambeck 1999. The greyscale indicates blue-shifted CO. Thick contours show SiO J=2-1 emission. Crosses of decreasing size give the positions of IRAS 2A, 2B, and 2C.

SiO in the E-W outflow but not the N-S, strongly suggesting that these are two separate outflows. Nonetheless, it still appears likely that the driving source of the N-S flow remains to be discovered, and further searches for binarity of IRAS 2A are suggested. In Figures 2 and 3 we present the higher resolution wideband and channel maps of SiO

in the E-W outflow. The much brighter, redshifted, eastern bow shock shows a wider range of velocities, and is associated with a dense clump seen in CS by Sandell *et al.*, and whose V-shaped geometry resolved by Blake (1996) suggests a bow shock driven into a steep density gradient. Presumably the eastern bow shock is much brighter because the driving jet is interacting with this dense clump. The actual emission peaks in both the east and west lobes are unresolved in our maps, and higher resolution observations are needed to confirm the expected bow shock geometry at the ends of the outflow lobes.



wilkin 31-Oct-2002 00:16

Figure 2. 3-Field mosaic of IRAS 2A outflow in SiO $j=2-1$. In this map, positions are relative to the bright, eastern emission peak. The submm point source locations are shown in Figures 1 and 3.




Figure 3. Channel maps at 5 km/s intervals. The large, medium and small crosses indicate the positions of IRAS 2A, 2B and 2C, respectively.

References

- Blake, G. A. 1996, IAU Symp. 178: Molecules in Astrophysics: Probes & Processes, 178, 31
- Chandler, C.J. & Richer, J.S. 2001, ApJ 555, 139
- Engargiola, G. & Plambeck, R. L. 1999, The Physics and Chemistry of the Interstellar Medium, Eds.: V. Ossenkopf, J. Stutzki, and G. Winnewisser
- Liseau, R., Sandell, G., Knee, L.B.G. 1988, A&A 192, 153
- Sandell, G., Knee, L.B.G., Aspin, C., Robson, I.E., & Russell, A.P.G. 1994, 1994, A&A 285, L1
- Sandell, G., Knee, L.B.G. 2001, ApJ 546, L49